

Student Illness Policy

A student should not be sent to school if he/she has the following symptoms or illnesses:

Temperature above 101 degrees axillary method. 100 degrees orally.
Diarrhea or vomiting
Chickenpox, scabies, ringworm, or impetigo
Head Lice
Symptoms of other contagious diseases (measles, hepatitis, scarlet fever, etc.)
A child does not feel well enough to participate in daily activities

Please consult the following chart to determine when your child may return to school:

Disease:	When child may return to school or your doctor's recommendation: (note must be presented)
Chicken Pox	Return to school six days after onset of blisters and lesions are crusted
Coxsackie (Hand, Foot & Mouth Disease)	Return to school when fever and sore throat have subsided for 24 hours .
Diarrhea	Return to school when diarrhea has subsided for 24 hours . Diarrhea is defined as six or more, loose, watery stools in 24 hours , e.g., a child has three loose stools at home and three at school should be excluded.
Fever	Return to school when fever has subsided for 24 hours without medication .
Head Lice	Return to school after first treatment has been completed AND the student is free of lice.
Impetigo/Staph/ MRSA	Return to school 24 hours after treatment with an antibiotic that has been started. The sores must be covered when the child returns to school.
Measles	Return to school on 5th day after appearance of rash.
Mumps	Return to school 10 days from onset of swelling.
Pink Eye (Conjunctivitis)	Return to school 24 hours after antibiotic treatment has been started
Ringworm	Return to school 24 hours after treatment has been started. The lesions must be covered (by clothing) when the child returns to school.
Rotavirus	Return to school when diarrhea has subsided for 24 hours .
RSV	Return to school when fever has subsided for 24 hours without use of medication.
Scabies	Return to school 24 hours after treatment with an appropriate medicated lotion.
Scarlet Fever	Return to school 24 hours after treatment with an antibiotic has been started.
Strep	Return to school 24 hours after treatment with an antibiotic and fever free for 24 hours without medication .
Vomiting	Return to school after vomiting has subsided for 24 hours .